

QCD Corrections in two-Higgs-doublet extensions of the Standard Model with Minimal Flavor Violation

G. Degrassi^a and P. Slavich^b

^a *Dipartimento di Fisica, Università di Roma Tre and INFN, Sezione di Roma Tre
Via della Vasca Navale 84, I-00146 Rome, Italy*

^b *LPTHE, 4, Place Jussieu, F-75252 Paris, France*

Abstract

We present the QCD corrections to R_b and to the $\Delta B = 1$ effective Hamiltonian in models with a second Higgs field that couples to the quarks respecting the criterion of Minimal Flavor Violation, thus belonging either to the $(\mathbf{1}, \mathbf{2})_{1/2}$ or to the $(\mathbf{8}, \mathbf{2})_{1/2}$ representation of $SU(3) \times SU(2) \times U(1)$. After the inclusion of the QCD corrections, the prediction for R_b becomes practically insensitive to the choice of renormalization scheme for the top mass, which for the type-I and type-II models translates in a more robust lower bound on $\tan \beta$. The QCD-corrected determinations of R_b and $\text{BR}(B \rightarrow X_s \gamma)$ are used to discuss the constraints on the couplings of a (colored) charged Higgs boson to top and bottom quarks.

e-mail:
degrassi@fis.uniroma3.it
slavich@lpthe.jussieu.fr

1 Introduction

One of the main goals of the present experimental program at the Tevatron and at the Large Hadron Collider (LHC) is the search for the Higgs boson(s) in order to elucidate the mechanism of electroweak symmetry breaking (EWSB). In the Standard Model (SM), the latter is realized in the most economical way via a single Higgs doublet. This minimal realization predicts a single neutral Higgs boson, whose mass can be constrained, from electroweak precision data and the direct search limit from LEP, to be lighter than ~ 200 GeV. However, at the moment, there is no direct experimental evidence for a neutral Higgs boson or any other scalar particle like, for example, a charged boson that can be present in models with a nonminimal Higgs sector.

The LHC is going to explore physics up to the TeV scale in order to search for the Higgs boson, as well as for any new phenomenon that would confirm the widespread expectation that the picture of particle physics in terms of the SM is incomplete. However, new particles with mass in the TeV range that couple to quarks at the tree level can modify the predictions for Flavor Changing Neutral Current (FCNC) processes. Thus any extension of the SM, starting from the simplest we can think of, a Two-Higgs-Doublet Model (2HDM), needs to face the problem of avoiding conflicts with the strict limits on FCNC processes.

Glashow and Weinberg addressed this issue proposing the principle of Natural Flavor Conservation (NFC) [1], which requires that the matrices of Yukawa couplings to up and down quarks for all the Higgs fields be diagonal in the basis where the quark mass matrices, $M^{U,D}$, are diagonal. This implies that, with the exception of models with vectorlike quarks that mix with the ordinary ones, NFC models do not have tree-level FCNC couplings. In the 2HDM case, NFC can be realized imposing the sufficient condition that each of the quark mass matrices is obtained from a single Higgs field. This can be enforced via a $\mathcal{Z}_2$ symmetry that acts differently on the two Higgs doublets, leading to two possibilities usually referred to as type-I (i.e., the model in which both up and down quarks get their masses from Yukawa couplings to the same Higgs doublet) and type-II models (where up and down quarks get their masses from Yukawa couplings to different Higgs doublets).

A less-restrictive way to suppress FCNC processes, still avoiding conflict with the experimental bounds, is to consider the criterion of Minimal Flavor Violation (MFV) [2], which amounts to assuming that all the new flavor-changing transitions, including those mediated at the tree level by electrically neutral particles, are controlled by the Cabibbo-Kobayashi-Maskawa (CKM) matrix. Thus, the MFV hypothesis requires that all the flavor-violating interactions of the new particles present at the TeV scale be linked to the known structure of the Yukawa couplings.

The enforcement of the MFV hypothesis to the case of multi-Higgs models has been recently investigated by several groups [3, 4, 5]. In particular, in ref. [3] it has been shown, via group-theoretic arguments, that the MFV hypothesis can be enforced requiring that all the Higgs Yukawa-coupling matrices be composed from the pair of matrices Y^U and Y^D that are responsible for the breaking of the $SU(3)_{Q_L} \times SU(3)_{U_R} \times SU(3)_{U_D}$ quark flavor symmetry. This requirement restricts the allowed $SU(3) \times SU(2) \times U(1)$ representations of the Higgs fields that can couple to the quarks to either be

equal to that of the SM Higgs field, i.e. $(\mathbf{1}, \mathbf{2})_{1/2}$, or transform as $(\mathbf{8}, \mathbf{2})_{1/2}$. Examples of the former case, besides the NFC type I and II models, are the aligned model of ref. [5] or the class of 2HDM presented in ref. [6]. The latter case is quite different, because the second field does not acquire a vacuum expectation value (vev) and does not mix with the SM Higgs field. Thus the scalar spectrum of this model contains a CP-even, color-singlet Higgs boson (the usual SM one) and three color-octet particles, one CP-even, one CP-odd and one electrically charged, which are split in mass proportionally to the SM-Higgs vev [3]. These colored scalar particles give rise to an interesting phenomenology for the LHC, not only because – if they are not too heavy – they can be directly produced, but also because their indirect effects can influence flavor, electroweak and Higgs physics [3, 7, 8, 9, 10].

Models with a second Higgs doublet present a new and interesting phenomenology, in particular related to the presence of a charged scalar. In the flavor sector, decays mediated by a weak charged current are the natural place where effects due to a charged Higgs boson, H^+ , can show up. In the electroweak sector the observable $R_b \equiv \Gamma(Z \rightarrow b\bar{b})/\Gamma(Z \rightarrow \text{hadrons})$ shows a sensitivity to H^+ because of the specific vertex corrections introduced by the interaction of H^+ with the top and bottom quarks. Many studies (for the most recent see, e.g., refs. [11, 12, 13]) used various combinations of flavor and electroweak observables to constrain the parameter space of the type-II 2HDM, which garnered most of the attention because of its property of having the same Higgs-sector realization as the Minimal Supersymmetric Standard Model (MSSM). Other studies [14, 15, 16] explored the parameter space of 2HDMs unconstrained by a $\mathcal{Z}_2$ symmetry, with the second Higgs doublet still in the $(\mathbf{1}, \mathbf{2})_{1/2}$ representation.

The theoretical accuracy of the predictions in the 2HDM with MFV is not yet at the same level as in the SM. Here we take a first step in improving this situation, by (re)considering the QCD corrections to two observables, R_b and $\text{BR}(B \rightarrow X_s \gamma)$, which allow to set important constraints on the mass and couplings of the charged scalar. The QCD corrections can play a relevant role in reducing the error of the theoretical predictions, a well-known example of this fact being indeed the radiative decay of the B meson. The present knowledge in the 2HDMs of the two observables we are considering can be summarized in this way: in the case of models with a second Higgs doublet in the $(\mathbf{1}, \mathbf{2})_{1/2}$ representation, the complete one-loop calculation of $\Gamma(Z \rightarrow b\bar{b})$ is available [17], but (to our knowledge) no QCD corrections to the charged-scalar contributions are known. The process $B \rightarrow X_s \gamma$ is instead fully known at the Next-to-Leading Order (NLO) in QCD [18, 19, 20, 21]. The case with colored scalars in the adjoint representation of $SU(3)$ is less studied. The one-loop charged-scalar contribution to $\Gamma(Z \rightarrow b\bar{b})$ was reported in ref. [7] (see also ref. [22]) while for the radiative decay of the B meson only a partial result for the Leading Order (LO) Wilson coefficients of the magnetic and chromo-magnetic operators has been presented [3].

In this paper we present the QCD corrections to the contribution to $\Gamma(Z \rightarrow b\bar{b})$ of a charged scalar in either the $(\mathbf{1}, \mathbf{2})_{1/2}$ or the $(\mathbf{8}, \mathbf{2})_{1/2}$ representation. Concerning $B \rightarrow X_s \gamma$, we compute the $\mathcal{O}(\alpha_s)$ contribution to the Wilson coefficients due to a colored charged scalar in the $(\mathbf{8}, \mathbf{2})_{1/2}$ representation. This is the missing piece to achieve NLO predictions for $\text{BR}(B \rightarrow X_s \gamma)$ for all 2HDMs with MFV.

Because of the specific interactions of the colored scalar with the gluons, the Wilson coefficients cannot be simply obtained by an appropriate color-factor rescaling of the known $(\mathbf{1}, \mathbf{2})_{1/2}$ result.

The paper is organized as follows: in the next section we discuss the couplings of the charged Higgs boson in the different realizations of the 2HDM with MFV. In section 3 we present the results for the QCD-corrected contribution to $\Gamma(Z \rightarrow b\bar{b})$ due to a charged scalar, covering both cases of color-singlet and color-octet particle. We show that, after the inclusion of the QCD correction, the prediction for R_b is practically insensitive to choice of an $\overline{\text{MS}}$ or on-shell (OS) renormalization scheme for the top mass. The bounds set by R_b on the tbH^+ coupling are also shown. Section 4 contains the result for the NLO Wilson coefficients in the $\Delta B = 1$ effective Hamiltonian (the explicit analytic expressions are presented in the appendix). The known results for the colorless 2HDMs are recovered, while the case of a colored charged scalar is fully new. The restrictions imposed by $B \rightarrow X_s \gamma$ on the charged-scalar interaction with quarks are discussed. Finally, in section 5 we present our conclusions.

2 Minimally flavor violating 2HDMs

In a generic 2HDM it is always possible to rotate the two Higgs fields to a basis in which only one of them, which we denote as Φ_1 , gets a vev [23]. In this basis, we write the Yukawa interactions of the Higgs fields with the quarks as

$$-\mathcal{L}_Y = \bar{q}_L \tilde{\Phi}_1 Y^U u_R + \bar{q}_L \Phi_1 Y^D d_R + \bar{q}_L \tilde{\Phi}_2^{(a)} T_R^{(a)} \bar{Y}^U u_R + \bar{q}_L \Phi_2^{(a)} T_R^{(a)} \bar{Y}^D d_R + \text{h.c.}, \quad (1)$$

where $\tilde{\Phi}_i \equiv i\sigma_2 \Phi_i^*$, and the Yukawa couplings $Y^{U,D}$ are 3×3 matrices in flavor space such that $M^{U,D} = Y^{U,D} \langle \Phi_1^0 \rangle$. The possibility of a colored second Higgs doublet is encoded in the matrices $T_R^{(a)}$ that act on the quark fields. In the usual colorless 2HDM T_R is equal to the identity matrix in color space. On the other hand, for a colored Higgs doublet in the adjoint representation of $SU(3)$ $T_R^a = T_F^a$ ($a = 1, 8$), the matrices of the fundamental representation. The MFV condition amounts to requiring that the Yukawa coupling matrices of the second doublet, $\bar{Y}^{U,D}$, be composed of combinations of the matrices $Y^{U,D}$, and transform under the $SU(3)_{Q_L} \times SU(3)_{U_R} \times SU(3)_{U_D}$ quark flavor symmetry in the same way as $Y^{U,D}$ themselves. We can decompose the matrices $\bar{Y}^{U,D}$ as

$$\bar{Y}^U = A_u \left(1 + \epsilon_u Y^U Y^{U\dagger} + \dots \right) Y^U, \quad \bar{Y}^D = A_d \left(1 + \epsilon_d Y^U Y^{U\dagger} + \dots \right) Y^D, \quad (2)$$

where in principle $A_{u,d}$ and $\epsilon_{u,d}$ are arbitrary complex coefficients. The ellipses in eq. (2) denote terms involving powers of $Y^D Y^{D\dagger}$ as well as terms involving higher powers of $Y^U Y^{U\dagger}$. In the following, we will assume that the only significant deviations from proportionality between $\bar{Y}^{U,D}$ and $Y^{U,D}$ are controlled by the Yukawa coupling of the top quark, and that terms involving higher powers of the Yukawa matrices are suppressed (e.g., because they are generated at higher loops). If we further require that there are no new sources of CP violation apart from the complex phase in the CKM matrix, the coefficients $A_{u,d}$ and $\epsilon_{u,d}$ must be real. Finally, the case $\epsilon_u = \epsilon_d = 0$ corresponds to the NFC situation in which the Yukawa matrices of both Higgs doublets are aligned in flavor space.

The processes $Z \rightarrow b\bar{b}$ and $B \rightarrow X_s \gamma$ that we will consider in sections 3 and 4 involve loops with a charged Higgs boson and a top quark. Under the assumptions implicit in eq. (2), the interaction between the quarks and H^+ is controlled by the Lagrangian

$$\mathcal{L}_{H^+} = -\frac{g}{\sqrt{2}m_W} \sum_{i,j=1}^3 \bar{u}_i T_R^{(a)} \left(A_u^i m_{u_i} \frac{1-\gamma_5}{2} - A_d^i m_{d_j} \frac{1+\gamma_5}{2} \right) V_{ij} d_j H_{(a)}^+ + \text{h.c.} , \quad (3)$$

where g is the $SU(2)$ coupling constant, i, j are generation indices, $m_{u,d}$ are quark masses, V is the CKM matrix. The family-dependent couplings $A_{u,d}^i$ read

$$A_{u,d}^i = A_{u,d} \left(1 + \epsilon_{u,d} \frac{m_t^2}{v^2} \delta_{i3} \right) , \quad (4)$$

where $v = \langle \Phi_1^0 \rangle$. It appears from eq. (4) that, when we neglect the masses of the light quarks, the effect on the charged-Higgs couplings arising from the $Y^U Y^{U\dagger}$ terms in eq. (2) is limited to a shift in the couplings involving the top quark. Since those are the only couplings that enter our computations, in sections 3 and 4 we will drop the family index from $A_{u,d}^i$ without ambiguity.

The term $\epsilon_d Y^U Y^{U\dagger}$ entering the expression for $\bar{Y}^D$ in eq. (2) also induces a flavor-changing interaction with the down quarks for the neutral component of the second Higgs doublet. However, this interaction does not affect the computation of $\Gamma(Z \rightarrow b\bar{b})$, and its contribution to $B \rightarrow X_s \gamma$ is negligible with respect to the charged-Higgs contribution as long as $\epsilon_d A_d/A_u \ll (v/m_b)^2$. Other FCNC processes such as $B\bar{B}$ mixing would put bounds on the combination $\epsilon_d A_d$, but this will not be relevant to the discussion that follows.

In the notation of eq. (3), the type-I and type-II models are specified by T_R^a equal to the identity matrix in color space, and by the real (and family-universal) coefficients

$$A_u^i = A_d^i = 1/\tan \beta \quad (\text{type I}), \quad (5)$$

$$A_u^i = -1/A_d^i = 1/\tan \beta \quad (\text{type II}), \quad (6)$$

where $\tan \beta$ is the ratio of the vevs of the two Higgs doublets in the basis where each of the quark mass matrices is obtained from a single Higgs field.

According to our discussion, the MFV hypothesis includes two other possibilities, namely color-singlet and color-octet Higgs doublet that couple to the quarks with arbitrary coefficients¹ $A_{u,d}^i$. We are going to refer to the first category (singlet) as type-III model, while the second possibility (octet) will be called type-C model.

¹Even models with generic Yukawa matrices not satisfying the MFV hypothesis show a structure of couplings with arbitrary coefficients $A_{u,d}^i$. To be phenomenologically viable, the dangerous FCNC effects should be sufficiently suppressed via some specific assumption like, e.g., a specific texture of the Yukawa matrices [24].

3 Charged-Higgs contribution to R_b including QCD corrections

We begin by discussing the radiatively corrected partial decay width of the Z boson in a quark-antiquark pair in a model with an additional Higgs doublet. We write it as

$$\Gamma(Z \rightarrow q\bar{q}) = N_c \frac{G_\mu}{\sqrt{2}} \frac{m_Z^3}{3\pi} [(\bar{g}_L^q)^2 + (\bar{g}_R^q)^2] K_q, \quad (7)$$

where N_c is the color factor ($N_c = 3$), $\bar{g}_{(L,R)}^q$ are the left-handed and right-handed $Zq\bar{q}$ couplings, written in terms of the radiative parameter ρ_q and the radiatively corrected sine of the Weinberg angle $\bar{s}_W^q$ as (T_3^q is the third component of the weak isospin, Q_q is the electric charge in unit e)

$$\begin{aligned} \bar{g}_L^q &= \sqrt{\rho_q} [T_3^q - Q_q (\bar{s}_W^q)^2], \\ \bar{g}_R^q &= -\sqrt{\rho_q} Q_q (\bar{s}_W^q)^2, \end{aligned} \quad (8)$$

while the factor K_q contains the QCD, QED and quark-mass corrections. The latter has been computed up to $\mathcal{O}(\alpha_s^3)$ in ref. [25], and at the lowest order it reads:

$$K_q = 1 + C_F \frac{3\alpha_s}{4\pi} + Q_q^2 \frac{3\alpha}{4\pi} - \frac{3}{4} \frac{\mu_q}{(\bar{g}_L^q)^2 + (\bar{g}_R^q)^2} + \mathcal{O}(\alpha_s^2), \quad (9)$$

where $\mu_q = m_q^2/m_Z^2$ and $C_F = 4/3$.

We assume that the oblique corrections due to the second doublet are negligible, as happens when the spectrum of the additional states is approximately custodially symmetric. Then the effect of the second Higgs doublet is concentrated in the vertex corrections to $\Gamma(Z \rightarrow b\bar{b})$. Thus, defining

$$\rho_q = \rho_q^{SM} + \delta\rho_q, \quad (10)$$

$$(\bar{s}_W^q)^2 = (\bar{s}_W^q)_{SM}^2 + \delta(\bar{s}_W^q)^2, \quad (11)$$

we have $\delta\rho_{(q \neq b)} = \delta(\bar{s}_W^{(q \neq b)})^2 = 0$. In the limit of neglecting the mass of the Z boson with respect to the masses of the top quark and the charged Higgs boson we find

$$\delta\rho_b = \frac{1}{T_3^b} \frac{\alpha}{4\pi s_W^2} C_R^1 \left[\left(\frac{|A_u| \hat{m}_t}{\sqrt{2} m_W} \right)^2 + \left(\frac{|A_d| \hat{m}_b}{\sqrt{2} m_W} \right)^2 \right] \left[f_1(t_h) + \frac{\alpha_s}{4\pi} (C_F f_2(t_h) + C_R^2 f_3(t_h)) \right], \quad (12)$$

$$\delta(\bar{s}_W^b)^2 = -\frac{1}{2} \delta\rho_b s_W^2 + \frac{1}{2 Q_b} \frac{\alpha}{4\pi s_W^2} C_R^1 \left(\frac{|A_d| \hat{m}_b}{\sqrt{2} m_W} \right)^2 \left[f_1(t_h) + \frac{\alpha_s}{4\pi} (C_F f_2(t_h) + C_R^2 f_3(t_h)) \right], \quad (13)$$

where $C_R^1 = 1, C_R^2 = 0$ [$C_R^1 = C_F, C_R^2 = C_A = N_c$] for Higgs fields in the $(\mathbf{1}, \mathbf{2})_{1/2}$ [$(\mathbf{8}, \mathbf{2})_{1/2}$] representation, and we omit an overall factor $|V_{tb}|^2 \approx 1$. In eqs. (12) and (13) $t_h = \hat{m}_t^2/m_{H^+}^2$, where $\hat{m}_q$ is the $\overline{\text{MS}}$ quark mass at the scale μ and m_{H^+} is the OS H^+ mass. The explicit expressions for the functions $f_i(x)$ are

$$f_1(x) = \frac{x}{x-1} - \frac{x \ln x}{(x-1)^2}, \quad (14)$$

$$\begin{aligned}
f_2(x) &= -\frac{6x(x-2)}{(x-1)^2} \text{Li}_2\left(1 - \frac{1}{x}\right) + \frac{x(-27+11x)}{(x-1)^2} + \frac{x(25-9x)\ln x}{(x-1)^3} \\
&+ \left(\frac{6x(3-x)}{(x-1)^2} - \frac{12x\ln x}{(x-1)^3}\right) \ln \frac{\hat{m}_t^2}{\mu^2} - 3f_1(x), \tag{15}
\end{aligned}$$

$$f_3(x) = \frac{3x}{(x-1)} \text{Li}_2\left(1 - \frac{1}{x}\right) + \frac{3x(1-2x+x^2+\ln^2 x)}{(x-1)^3} - \frac{6x\ln x}{(x-1)^2}, \tag{16}$$

where the last term ($-3f_1$) in the function f_2 is introduced to avoid double counting due to the correction factor K_q in eq. (7), and in the tbH^+ coupling we have also kept the contribution proportional to the bottom mass².

The one-loop terms in $\delta\rho_b$ and $\delta(\bar{s}_W^b)^2$ agree with the results³ of refs. [17, 22]. The two-loop terms were obtained following the lines of the analogous SM calculation that was performed by several groups, via different methods, in the early nineties [26]. The SM correction can be actually obtained by considering the SM Lagrangian in the limit of vanishing gauge coupling constants, the so-called gaugeless limit of the SM [27]. In this limit the gauge bosons play the role of external sources, and the propagating fields are those of a Yukawa theory with massless Goldstone bosons. Indeed, in the limit $m_{H^+} \rightarrow 0$, the $\mathcal{O}(\alpha\alpha_s\hat{m}_t^2/m_W^2)$ corrections in eq. (12) agree with the known SM result.

To express the corrections in terms of the OS top mass m_t , we must expand the $\overline{\text{MS}}$ top mass entering the one-loop part as $\hat{m}_t = m_t + \delta m_t$, with

$$\delta m_t = \frac{\alpha_s}{4\pi} C_F \left(3 \ln \frac{m_t^2}{\mu^2} - 4\right) m_t. \tag{17}$$

For the terms proportional to $|A_u|^2$ and $|A_d|^2$ in eqs. (12) and (13), this amounts to making the substitution $\hat{m}_t \rightarrow m_t$ and replacing the function f_2 with the OS counterparts f_2^u and f_2^d , respectively:

$$f_2^u(x) = f_2(x) + \frac{8\pi}{\alpha_s C_F} \frac{\delta m_t}{m_t} \left[f_1(x) + x \frac{\partial f_1(x)}{\partial x} \right], \quad f_2^d(x) = f_2(x) + \frac{8\pi}{\alpha_s C_F} \frac{\delta m_t}{m_t} x \frac{\partial f_1(x)}{\partial x}. \tag{18}$$

The explicit dependence on the renormalization scale cancels out in the function f_2^u :

$$f_2^u(x) = -\frac{6x(x-2)}{(x-1)^2} \text{Li}_2\left(1 - \frac{1}{x}\right) + \frac{3x}{(x-1)} - \frac{9x\ln x}{(x-1)^2} - 3f_1(x), \tag{19}$$

whereas f_2^d has a residual dependence on μ which is compensated for by the implicit scale dependence of the $\hat{m}_b^2$ entering the one-loop parts of eqs. (12) and (13). Using the OS bottom mass m_b would remove this residual scale dependence, but it would introduce large logarithms of the ratio m_t/m_b in the two-loop part of the corrections.

In fig. 1 we plot, as a function of m_{H^+} , the ratio δ^{QCD} between the two-loop and one-loop contributions in the terms proportional to $|A_u|^2$ in eq. (12), for the two cases of color-singlet (lower set of lines) and color-octet (upper set of lines) charged Higgs boson. For each case, we show δ^{QCD} as

²Terms proportional to $\hat{m}_b$, relevant only for very large values of A_d , can also arise from vertices with neutral scalars.

³In ref. [22] there is a misprint in the overall normalization of the $\delta g^{L,R}$ couplings.




Figure 1: the ratio of two-loop to one-loop charged-Higgs contributions to the $Zb\bar{b}$ vertex as a function of m_{H^+} , with the top mass expressed in the $\overline{\text{MS}}$ (solid lines) or OS (dashed lines) renormalization scheme. The upper (red) curves are for the model with color-octet Higgs (type C), while the lower (blue) curves are for the models with color-singlet Higgs (types I–III).

obtained using either the central value of the physical (OS) top mass, $m_t = 173.1$ GeV [28] (dashed lines), or the corresponding $\overline{\text{MS}}$ value $\hat{m}_t(m_t) = 163.5$ GeV (solid lines), with the appropriate formulae for the two-loop function f_2 . It can be seen from the figure that, for models of types I–III (i.e. with color-singlet charged Higgs) the two-loop corrections are always negative, and they are substantially larger when the OS top mass is used in the one-loop part than when the $\overline{\text{MS}}$ mass is used. On the other hand, for the model of type C (with color-octet charged Higgs) there is an overall upward shift in the two-loop correction due to the additional function f_3 in eq. (12), with the result that the sign and relative size of the corrections in the OS and $\overline{\text{MS}}$ cases depend on the Higgs mass. For low values $m_{H^+} \approx 150$ GeV, the two-loop correction approaches zero if the OS top mass is used, and it is positive and relatively large if the $\overline{\text{MS}}$ mass is used. Conversely, for larger values $m_{H^+} \approx 400$ GeV the two-loop correction approaches zero if the $\overline{\text{MS}}$ top mass is used, and it is negative and relatively large if the OS mass is used. As a result, we will see that for 2HDMs of types I–III a reliable approximation of the two-loop result for R_b could be obtained by using the one-loop result expressed in terms of the $\overline{\text{MS}}$ top mass. On the other hand, a precise determination of R_b in the 2HDM of type C requires the inclusion of the two-loop part of the $Zb\bar{b}$ vertex correction.

From eqs. (7)–(13) we can construct the observable R_b , which can be written as

$$\frac{1}{R_b} = 1 + \frac{\sum_{(q \neq b)} [(\bar{g}_L^q)^2 + (\bar{g}_R^q)^2] K_q}{[(\bar{g}_L^b)^2 + (\bar{g}_R^b)^2] K_b} \equiv 1 + \frac{S_b}{s_b} C_b \quad (20)$$

where

$$S_b = \sum_{(q \neq b)} s_q ; \quad s_q = [(\bar{g}_L^q)^2 + (\bar{g}_R^q)^2] \left(1 + Q_q^2 \frac{3\alpha}{4\pi} \right) . \quad (21)$$

Using the results of ref. [29] we find $S_b = 0.6607$. Concerning the SM part of s_b , to avoid relying indirectly on the measured value of R_b , we follow ref. [12] and compute it using the values $\rho_b^{SM} = 0.99426$ [29] and $(\bar{s}_W^b)_{SM}^2 = (1.0063) \times 0.23153 = 0.23299$, the latter obtained from the measured value of $\sin^2 \theta_{eff}^{lept} = 0.23153 \pm 0.00016$ corrected for the top-induced contributions specific to the $Zb\bar{b}$ vertex⁴. Finally the factor C_b that includes QCD and mass corrections is obtained from ref. [25]. We find, for $\alpha_s = 0.118$, $C_b = 1.0086$.

With the values specified above for the various parameters entering eqs. (20) and (21) we find a SM prediction $R_b^{SM} = 0.21580$, nearly 1σ below the measured value $R_b^{\text{exp}} = 0.21629 \pm 0.00066$ [29]. Since the charged-Higgs contributions to $\bar{g}_L^b$ and $\bar{g}_R^b$ have the effect of further lowering R_b , stringent bounds can be imposed on the parameters m_{H^+} and A_u by the requirement that the predicted value of R_b in a 2HDM be not too far from R_b^{exp} . On the other hand, R_b has little sensitivity on A_d , because the terms in eqs. (12) and (13) controlled by the latter are suppressed by $\hat{m}_b$. Therefore, for the models of types III and C in which A_d is a free parameter we will simplify our discussion by setting $A_d = 0$. In the 2HDMs of type I and II the parameter A_d is related to A_u and cannot be set independently to zero. However, due to the strong suppression of the contributions controlled by A_d , in most of the parameter space the predictions of R_b obtained in those two models do not differ significantly from the predictions obtained in the type-III 2HDM with $A_d = 0$. More specifically, the predictions of the type-I 2HDM, in which $A_d = A_u$, are virtually indistinguishable from those of the type-III 2HDM with $A_d = 0$ for all the values of A_u consistent with the measured value of R_b^{exp} . In the 2HDM of type II, on the other hand, $A_d = -1/A_u$, and the predictions of R_b differ from the ones obtained in the type-III 2HDM with $A_d = 0$ only for very small values of A_u .

Figures 2 and 3 show our determination of R_b in the 2HDMs of type III and C, respectively, as a function of $|A_u|$. In each plot we show two sets of curves for the charged-Higgs mass values $m_{H^+} = 100$ GeV and $m_{H^+} = 400$ GeV. In each set, the dashed (solid) curve represents the one-loop (two-loop) result expressed in terms of the $\overline{\text{MS}}$ top mass, while the dotted (dot-dashed) curve represents the one-loop (two-loop) result expressed in terms of the physical top mass. We also show in each plot the measured value R_b^{exp} (solid horizontal line) and the values 1σ and 2σ below (dashed horizontal lines). It can be seen that, in both plots, the curves corresponding to the two-loop results (with the top mass renormalized either in the $\overline{\text{MS}}$ or in the OS scheme) are practically overlapped. The location of the curves corresponding to the one-loop results reflects the behavior that could already be inferred from

⁴ In ref. [12] the correcting factor 1.0063 was not introduced.




Figure 2: R_b as a function of $|A_u|$ in the 2HDM with color-singlet Higgs, for $A_d = 0$ and two different values of m_{H^+} . The measured value $R_b^{\text{exp}} = 0.21629$ and the values 1σ and 2σ below it are displayed as horizontal lines. For the meaning of the different curves see the text.




Figure 3: same as figure 2 in the 2HDM with color-octet Higgs.

fig. 1: in the 2HDM of type III the one-loop result computed in terms of the $\overline{\text{MS}}$ top mass is very close to the two-loop result, while the one-loop result computed in terms of the physical top mass can differ significantly. On the other hand, in the 2HDM of type C the quality of the one-loop approximation depends on the charged Higgs mass. At low values of m_{H^+} , using the physical top mass in the one-loop result gives a much better approximation to the two-loop result than using the $\overline{\text{MS}}$ top mass, while the situation is reversed at large values of m_{H^+} . Therefore, only the use of the two-loop results guarantees a precise determination of R_b for all the values of m_{H^+} .

From figures 2 and 3 it is also possible to determine the values of $|A_u|$ that are disfavored by the comparison between R_b^{exp} and the corresponding theoretical prediction. In the case of the type-III 2HDM, the two-loop curves cross the 2σ horizontal line at $|A_u| = 0.78$ for $m_{H^+} = 100$ GeV and at $|A_u| = 1.35$ for $m_{H^+} = 400$ GeV. In the case of the type-C 2HDM, they cross it at $|A_u| = 0.62$ for $m_{H^+} = 100$ GeV and at $|A_u| = 1.10$ for $m_{H^+} = 400$ GeV. We checked that the crossing points for intermediate values of m_{H^+} can be determined by linear interpolation of the values given above.

The predictions of R_b for the type-II 2HDM (where $A_d = -1/A_u = -\tan\beta$) are virtually indistinguishable from those presented in figure 2 for the type-III 2HDM as soon as $|A_u| > 0.1$. The upper bounds on $|A_u|$ discussed above translate, for the type-II 2HDM, into $|A_d| > 1.28$ for $m_{H^+} = 100$ GeV and $|A_d| > 0.74$ for $m_{H^+} = 400$ GeV. On the other hand, when $|A_u|$ tends to zero the predictions of R_b in the type-II 2HDM decrease quickly, and get two standard deviations below R_b^{exp} for $|A_u| \approx 0.01$, corresponding to $|A_d| \approx 100$. As we will see in the next section, the bounds on A_d coming from the process $B \rightarrow X_s \gamma$ can be much stronger than that, but they do not apply to the type-II 2HDM.

4 Charged-Higgs contribution to $B \rightarrow X_s \gamma$ at the NLO

The branching ratio for $B \rightarrow X_s \gamma$ is fully known at the NLO for a 2HDM of types I–III. To cover also the case of a type-C 2HDM, the only missing ingredient is the determination of the colored-scalar $\mathcal{O}(\alpha_s)$ contribution to the Wilson coefficients. We compute it following the analogous computation for the type I–II 2HDM presented in ref. [19].

In the operator basis defined in ref. [19] we write the Wilson coefficients at the scale μ_W , where the “full” theory is matched to an effective theory with five quark flavors, as

$$C_i(\mu_W) = C_i^{(0)}(\mu_W) + \delta C_i^{(0)}(\mu_W) + \frac{\alpha_s(\mu_W)}{4\pi} \left[C_i^{(1)}(\mu_W) + \delta C_i^{(1)}(\mu_W) \right], \quad (22)$$

where $C_i^{(k)}(\mu_W)$ represents the SM contribution ($k = 0, 1$) while $\delta C_i^{(k)}(\mu_W)$ represents the charged-Higgs contribution. At the LO, the latter is given by

$$\delta C_i^{(0)}(\mu_W) = 0 \quad i = 1, \dots, 6, \quad (23)$$

$$\delta C_7^{(0)}(\mu_W) = C_R^1 \left(\frac{|A_u|^2}{3} F_7^{(1)}(y) - A_d A_u^* F_7^{(2)}(y) \right), \quad (24)$$

$$\delta C_8^{(0)}(\mu_W) = C_R^1 \left(\frac{|A_u|^2}{3} F_8^{(1)}(y) - A_d A_u^* F_8^{(2)}(y) \right) + C_R^2 \left(|A_u|^2 F_8^{(3)}(y) + A_d A_u^* F_8^{(4)}(y) \right), \quad (25)$$

where

$$F_7^{(1)}(y) = \frac{y(7-5y-8y^2)}{24(y-1)^3} + \frac{y^2(3y-2)}{4(y-1)^4} \ln y, \quad F_7^{(2)}(y) = \frac{y(3-5y)}{12(y-1)^2} + \frac{y(3y-2)}{6(y-1)^3} \ln y, \quad (26)$$

$$F_8^{(1)}(y) = \frac{y(2+5y-y^2)}{8(y-1)^3} - \frac{3y^2}{4(y-1)^4} \ln y, \quad F_8^{(2)}(y) = \frac{y(3-y)}{4(y-1)^2} - \frac{y}{2(y-1)^3} \ln y, \quad (27)$$

$$F_8^{(3)}(y) = \frac{y(1+y)}{16(y-1)^2} - \frac{y^2}{8(y-1)^3} \ln y, \quad F_8^{(4)}(y) = -\frac{y}{4(y-1)} + \frac{y}{4(y-1)^2} \ln y, \quad (28)$$

with

$$y = \frac{\hat{m}_t^2(\mu_W)}{m_{H^+}^2}, \quad (29)$$

expressed in terms of the NLO top-quark running mass at the scale μ_W and of the OS charged-Higgs mass. Again, $C_R^1 = 1, C_R^2 = 0$ [$C_R^1 = C_F, C_R^2 = C_A = N_c$] for Higgs fields in the $(\mathbf{1}, \mathbf{2})_{1/2}$ [$(\mathbf{8}, \mathbf{2})_{1/2}$] representation.

At the NLO, the charged-Higgs contributions to the Wilson coefficients are

$$\delta C_i^{(1)}(\mu_W) = 0 \quad i = 1, 2, 3, 5, 6, \quad (30)$$

$$\delta C_4^{(1)}(\mu_W) = E^H(y), \quad (31)$$

$$\delta C_7^{(1)}(\mu_W) = G_7^H(y) + \Delta_7^H(y) \ln \frac{\mu_W^2}{m_{H^+}^2}, \quad (32)$$

$$\delta C_8^{(1)}(\mu_W) = G_8^H(y) + \Delta_8^H(y) \ln \frac{\mu_W^2}{m_{H^+}^2}. \quad (33)$$

The expressions for $G_{7,8}^H$, $\Delta_{7,8}^H$ and E^H are rather long and they are reported in the appendix. As expected, the μ_W dependence in $\delta C_{7,8}$ cancels out at $\mathcal{O}(\alpha_s)$ because the functions $\Delta_{7,8}^H$ entering eqs. (32) and (33) satisfy the relation

$$\Delta_i^H = \gamma_0^m y \frac{\partial \delta C_i^{(0)}}{\partial y} + \frac{1}{2} \sum_{j=1}^8 \gamma_{ji}^{(0)eff} \delta C_j^{(0)}, \quad (34)$$

where $\gamma_0^m = 8$ is the LO anomalous dimension of the top mass, while $\gamma_{ji}^{(0)eff}$ is the matrix of LO anomalous dimensions of the Wilson coefficients, whose entries can be found in eq. (8) of ref. [18].

In fig. 4 we show the ratio between the NLO and LO charged-Higgs contributions to the Wilson coefficients $C_{7,8}$ as a function of m_{H^+} , for both the type-C (color-octet) and type-III (color-singlet) cases, with the particular choice $A_d = A_u$ (in the color-singlet case this coincides with the type-I 2HDM). For the type-C 2HDM, the NLO corrections can reach up to $\sim 20\%$ of the LO contribution at small m_{H^+} , and they decrease as the charged-Higgs mass increases, eventually crossing zero. In contrast, for the type-III 2HDM the two-loop corrections to the Wilson coefficients are always different from zero, and have opposite sign with respect to the LO contributions. We checked that the behavior of the NLO corrections is qualitatively similar to the one described above even when we allow A_d to take on values different from A_u .




Figure 4: the ratio of NLO to LO charged-Higgs contributions to the Wilson coefficients C_7 (solid line) and C_8 (dashed line) as a function of m_{H^+} , with $A_d = A_u$, for the model with color-octet Higgs (type C) or color-singlet Higgs (type III).

The calculation of $\text{BR}(B \rightarrow X_s \gamma)$ is performed using a modified version of the fortran code **SusyBSG** [30]. The code provides a NLO evaluation of the branching ratio in the MSSM with MFV, including the full two-loop gluino contributions to the Wilson coefficients [31]. The current public version (1.3) includes also the options of evaluating the branching ratio in the SM, in the type-II 2HDM and in the MSSM with two-loop gluino contributions computed in the effective Lagrangian approximation. We enlarged the 2HDM option to include also the type-I, type-III and type-C models, thus covering all four types of 2HDM compatible with MFV.⁵ The relation between the Wilson coefficients and $\text{BR}(B \rightarrow X_s \gamma)$ is computed at NLO along the lines of ref. [32], but the free renormalization scales entering the NLO calculation are adjusted in such a way as to mimic the Next-to-Next-to-Leading Order (NNLO) contributions presented in ref. [33]. When the SM input parameters are set to the partially outdated values used in ref. [33], **SusyBSG** gives a SM prediction for $\text{BR}(B \rightarrow X_s \gamma)$ of 3.15×10^{-4} , in full agreement with the NNLO result of that paper. Very good numerical agreement is also found with the results of the partial NNLO implementation of the type-II 2HDM in ref. [33], which combines NNLO anomalous dimensions and matrix elements with NLO Wilson coefficients. We

⁵A public version of **SusyBSG** with this new feature will be released soon.




Figure 5: $\text{BR}(B \rightarrow X_s \gamma)$ as a function of A_d in the type-III (solid line) and type-C (dashed line) models for $A_u = 0.3$ (left panel) and $A_u = -0.3$ (right panel) and two different values of m_{H^+} . The horizontal dashed lines specify the 2σ interval around the experimental value of $\text{BR}(B \rightarrow X_s \gamma)$.

take into account a recent update [34] in the calculation of the normalization factor for the branching ratio as well as the latest central value of the top mass [28], which results in a modest enhancement of the SM prediction for $\text{BR}(B \rightarrow X_s \gamma)$ to 3.28×10^{-4} .

In the 2HDMs of types I and II, the requirement of consistency between the theoretical prediction and the measured value of $\text{BR}(B \rightarrow X_s \gamma)$ allows us to set bounds on the parameters m_{H^+} and $\tan \beta$, the latter determining both Higgs-quark couplings A_u and A_d . More specifically, in the type-I 2HDM the charged-Higgs contribution to the Wilson coefficients $C_{7,8}$ scales like $1/\tan \beta^2$, therefore it is possible to derive, for each given value of m_{H^+} , a lower bound on $\tan \beta$ (i.e., an upper bound on $A_u = A_d$) which is however much less stringent than the corresponding bound derived from R_b . In the type-II 2HDM there is a $\tan \beta$ -independent contribution to the Wilson coefficient, which allows to set an absolute lower bound on m_{H^+} . These bounds have been extensively discussed in the literature (see, e.g., refs. [11, 12, 13]) and we will not further consider them here. In the models of type III and C, on the other hand, the parameters A_u and A_d are unrelated to each other. As can be seen in eqs. (24) and (25), the charged-Higgs contributions to the Wilson coefficients $C_{7,8}$ include two terms controlled by $|A_u|^2$ and $A_d A_u^*$, respectively. The bounds on A_u derived in the previous section tell us that the former term cannot be too large, while the latter can be significant for large values of A_d . Furthermore, its effect on the branching ratio depends on the relative sign between A_u and A_d .

In fig. 5 we show $\text{BR}(B \rightarrow X_s \gamma)$ as a function of A_d , for both the type-III (solid lines) and type-C (dashed lines) 2HDM, and for the representative choices $A_u = \pm 0.3$ (the latter are allowed by R_b , as can be seen in figs. 2 and 3). The left panel displays the case of same sign between A_u and A_d , while the right panel shows the case of opposite sign. Each plot contains two sets of curves corresponding




Figure 6: $\text{BR}(B \rightarrow X_s \gamma)$ as a function of A_d in the type-III (solid line) and type-C (dashed line) models for $A_u = 0.3$ and two different values of m_{H^+} . The horizontal dashed lines specify the 2σ interval around the experimental value of $\text{BR}(B \rightarrow X_s \gamma)$.

to $m_{H^+} = 100$ GeV and $m_{H^+} = 400$ GeV, respectively. The horizontal dashed lines mark the 95% C.L. band around the experimental value $\text{BR}(B \rightarrow X_s \gamma) = (3.52 \pm 0.25) \times 10^{-4}$ [35]. The band also includes, added in quadrature, the theoretical error on the 2HDM prediction (we conservatively estimate this error as 10% of the SM prediction for the branching ratio). From fig. 5 it is clear that, unless $|A_u|$ is extremely small, the process $B \rightarrow X_s \gamma$ sets stringent limits on A_d . Focusing on the case of color-singlet Higgs we see that, for $A_u = 0.3$, the values of A_d that allow for a branching ratio inside the 95% C.L. band are $A_d \leq 0.9$ for $m_{H^+} = 100$ GeV and $A_d \leq 2.5$ for $m_{H^+} = 400$ GeV; for $A_u = -0.3$ the bounds are a little less stringent, i.e. $A_d \leq 1.3$ for $m_{H^+} = 100$ GeV and $A_d \leq 4$ for $m_{H^+} = 400$ GeV. From the figure it is also apparent that the bound on A_d for a fixed value of A_u is almost independent of the colored or colorless nature of the charged Higgs, with the colored case showing only slightly stronger bounds. However, as seen in the previous section, the bounds on A_u derived from R_b are more dependent on the nature of the Higgs, so that the allowed regions for the A_d coefficient are in fact different for the color-singlet and color-octet charged Higgs.

The case of same sign between A_u and A_d has the peculiarity that, as shown in fig. 6, there are actually two ranges of values of A_d that fit inside the 2σ allowed band for $\text{BR}(B \rightarrow X_s \gamma)$. This is related to the fact that, in this case, the sign of the charged-Higgs contribution to the Wilson coefficient of the magnetic dipole operator, C_7^H , is opposite to the sign of the SM contribution, C_7^{SM} . Since $\text{BR}(B \rightarrow X_s \gamma)$ is roughly proportional to $|C_7^H + C_7^{\text{SM}}|^2$, as A_d increases the branching ratio goes to zero when $C_7^H \approx -C_7^{\text{SM}}$, and it goes back inside the 2σ -allowed band when $C_7^H \approx -2C_7^{\text{SM}}$. Thus,

the two ranges of possible values of A_d differ by the sign of the amplitude $\mathcal{A}(b \rightarrow s\gamma)$, basically the sign of the Wilson coefficient C_7 . Although $B \rightarrow X_s \gamma$ allows both ranges of values for A_d , there are other observables that are sensitive to the sign of C_7 , thus selecting one of the two options. Among them, we cite $\text{BR}(B \rightarrow X_s l^+ l^-)$ [36] and the isospin-breaking asymmetry that can be constructed from the exclusive neutral and charged $B \rightarrow K^* \gamma$ decay modes [37, 16]. Although, for both observables, neither the experimental result nor the theoretical prediction is at the same level of accuracy as for $\text{BR}(B \rightarrow X_s \gamma)$, these observables still give a compelling indication that the sign of $\mathcal{A}(b \rightarrow s\gamma)$ is that of the SM contribution, thus eliminating the large- A_d option.

5 Conclusions

Minimal flavor violation is a very popular criterion that is used to suppress FCNC effects in models with new particles at the TeV scale. The enforcement of the MFV criterion to the simplest extension of the SM, i.e. a model with a second Higgs doublet, allows the possibility of color-singlet or color-octet Higgs field. For both cases we have considered the two-loop QCD corrections to the charged Higgs boson contribution to R_b . We found that for all four types of 2HDM with MFV, after the inclusion of the two-loop QCD corrections, the prediction for this observable is practically insensitive to the choice of renormalization scheme for the top mass entering the one-loop part of the calculation. Thus, the upper bound on the coupling $|A_u|$ derived from R_b is improved, which for the type-I and type-II models translates in a more robust lower bound on $\tan\beta$. We have also computed the $\mathcal{O}(\alpha_s)$ contributions to the Wilson coefficients relevant to the process $B \rightarrow X_s \gamma$ for the color-octet case. This was the last missing ingredient to obtain a determination of $\text{BR}(B \rightarrow X_s \gamma)$ at the NLO level for all 2HDM with MFV. After the inclusion of the NLO corrections it is found that, in the region allowed by the present experimental results, the $B \rightarrow X_s \gamma$ transition is fairly insensitive to the colored or colorless nature of the charged Higgs. Furthermore, in type-III and C models, the bounds on the A_d parameter that can be obtained from $\text{BR}(B \rightarrow X_s \gamma)$ and other observables rule out the large- A_d region, where effects proportional to the bottom mass could become important.

Acknowledgments

We thank H. Haber for communications concerning ref. [22], and P. Gambino for useful discussions. One of us (G.D.) also thanks M. Nebot for his contribution in the early stage of this project. This work was supported in part by an EU Marie-Curie Research Training Network under contract MRTN-CT-2006-035505 (HEPTOOLS) and by ANR under contract BLAN07-2_194882.

Appendix: Analytical expressions for the NLO Wilson coefficients

In this appendix we report the analytic expressions for the functions $G_{7,8}^H$ and $\Delta_{7,8}^H$ entering $\delta C_{7,8}^{(1)}$. We find

$$\begin{aligned}
G_7^H(y) = & C_R^1 C_F \left\{ A_d A_u^* y \left[\frac{4(-3+7y-2y^2)}{3(y-1)^3} \text{Li}_2 \left(1 - \frac{1}{y} \right) + \frac{8-14y-3y^2}{3(y-1)^4} \ln^2 y \right. \right. \\
& \left. \left. + \frac{2(-3-y+12y^2-2y^3)}{3(y-1)^4} \ln y + \frac{7-13y+2y^2}{(y-1)^3} \right] \right. \\
& + |A_u|^2 y \left[\frac{y(18-37y+8y^2)}{6(y-1)^4} \text{Li}_2 \left(1 - \frac{1}{y} \right) + \frac{y(-14+23y+3y^2)}{6(y-1)^5} \ln^2 y \right. \\
& \left. + \frac{-50+251y-174y^2-192y^3+21y^4}{54(y-1)^5} \ln y \right. \\
& \left. \left. + \frac{797-5436y+7569y^2-1202y^3}{648(y-1)^4} \right] \right\} \\
& + C_R^1 C_R^2 \left\{ A_d A_u^* y \left[\frac{-19+25y}{18(y-1)^2} \text{Li}_2 \left(1 - \frac{1}{y} \right) + \frac{-25+33y+15y^2}{36(y-1)^4} \ln^2 y \right. \right. \\
& \left. \left. + \frac{33-59y+3y^2}{18(y-1)^3} \ln y + \frac{-8+31y}{36(y-1)^2} \right] \right. \\
& + |A_u|^2 y \left[\frac{12-25y}{36(y-1)^2} \text{Li}_2 \left(1 - \frac{1}{y} \right) + \frac{-12+85y-108y^2+3y^3}{72(y-1)^5} \ln^2 y \right. \\
& \left. \left. - \frac{50+33y-195y^2+16y^3}{108(y-1)^4} \ln y + \frac{17-29y+4y^2}{18(y-1)^3} \right] \right\} , \quad (A1)
\end{aligned}$$

$$\begin{aligned}
\Delta_7^H(y) = & C_R^1 C_F \left\{ A_d A_u^* y \left[\frac{21-47y+8y^2}{6(y-1)^3} + \frac{-8+14y+3y^2}{3(y-1)^4} \ln y \right] \right. \\
& \left. + |A_u|^2 y \left[\frac{-31-18y+135y^2-14y^3}{36(y-1)^4} + \frac{y(14-23y-3y^2)}{6(y-1)^5} \ln y \right] \right\} \\
& + C_R^1 C_R^2 \left\{ A_d A_u^* y \left[\frac{1}{3(y-1)} - \frac{1}{3(y-1)^2} \ln y \right] \right. \\
& \left. + |A_u|^2 y \left[-\frac{1+y}{12(y-1)^2} + \frac{y}{6(y-1)^3} \ln y \right] \right\} , \quad (A2)
\end{aligned}$$

$$\begin{aligned}
G_8^H(y) = & C_R^1 \left\{ A_d A_u^* \frac{1}{3} y \left[\frac{-36+25y-17y^2}{2(y-1)^3} \text{Li}_2 \left(1 - \frac{1}{y} \right) + \frac{19+17y}{(y-1)^4} \ln^2 y \right. \right. \\
& \left. \left. + \frac{-3-187y+12y^2-14y^3}{4(y-1)^4} \ln y + \frac{3(143-44y+29y^2)}{8(y-1)^3} \right] \right. \\
& + |A_u|^2 \frac{1}{6} y \left[\frac{y(30-17y+13y^2)}{(y-1)^4} \text{Li}_2 \left(1 - \frac{1}{y} \right) - \frac{y(31+17y)}{(y-1)^5} \ln^2 y \right. \\
& \left. \left. + \frac{-226+817y+1353y^2+318y^3+42y^4}{36(y-1)^5} \ln y \right] \right\}
\end{aligned}$$

$$\begin{aligned}
& + \frac{1130 - 18153y + 7650y^2 - 4451y^3}{216(y-1)^4} \Big] \Big\} \\
& + C_R^2 \left\{ A_d A_u^* \frac{1}{9} y \left[\frac{-43 + 34y}{4(y-1)^2} \text{Li}_2 \left(1 - \frac{1}{y} \right) + \frac{-157 - 108y + 81y^2}{8(y-1)^4} \ln^2 y \right. \right. \\
& \quad \left. \left. + \frac{-51 - 22y + 57y^2}{8(y-1)^3} \ln y + \frac{5(13 - 8y)}{(y-1)^2} \right] \right. \\
& \quad + |A_u|^2 \frac{1}{144} y \left[\frac{-15 + 149y - 122y^2}{(y-1)^3} \text{Li}_2 \left(1 - \frac{1}{y} \right) \right. \\
& \quad \left. - \frac{15 - 533y - 237y^2 + 243y^3}{2(y-1)^5} \ln^2 y \right. \\
& \quad \left. \left. - \frac{172 - 744y + 357y^2 + 23y^3}{3(y-1)^4} \ln y - \frac{203 + 1174y - 737y^2}{2(y-1)^3} \right] \right\} \quad (\text{A3})
\end{aligned}$$

$$\begin{aligned}
\Delta_8^H(y) &= C_R^1 \left\{ A_d A_u^* \frac{1}{3} y \left[\frac{81 - 16y + 7y^2}{2(y-1)^3} - \frac{19 + 17y}{(y-1)^4} \ln y \right] \right. \\
&\quad \left. + |A_u|^2 \frac{1}{6} y \left[\frac{-38 - 261y + 18y^2 - 7y^3}{6(y-1)^4} + \frac{y(31 + 17y)}{(y-1)^5} \ln y \right] \right\} \\
&+ C_R^2 \left\{ A_d A_u^* \frac{1}{6} y \left[\frac{31 - 7y}{(y-1)^2} - \frac{19 + 5y}{(y-1)^3} \ln y \right] \right. \\
&\quad \left. + |A_u|^2 \frac{1}{12} y \left[\frac{-19 - 60y + 7y^2}{2(y-1)^3} + \frac{y(31 + 5y)}{(y-1)^4} \ln y \right] \right\} \quad (\text{A4})
\end{aligned}$$

$$\begin{aligned}
E^H(y) &= C_R^1 |A_u|^2 \frac{1}{6} y \left[\frac{16 - 29y + 7y^2}{6(y-1)^3} + \frac{-2 + 3y}{(y-1)^4} \ln y \right] \\
&+ C_R^2 |A_u|^2 \frac{1}{4} y \left[\frac{-1}{(y-1)} + \frac{2 + y}{3(y-1)^2} \ln y \right] . \quad (\text{A5})
\end{aligned}$$

The results for type I-III models are recovered setting $C_R^1 = 1$ and $C_R^2 = 0$, while the result for the case of a colored charged scalar in the adjoint of $SU(3)$ is obtained with $C_R^1 = C_F$ and $C_R^2 = N_c$.

Note added: after the publication of our paper, we became aware of ref. [38], which contains two-loop formulae for the Wilson coefficients of the operators relevant to $b \rightarrow s\gamma$ in a generic model with a heavy fermion and a heavy scalar. While ref. [38] does not specifically discuss the case of the type-C 2HDM, the results presented in our appendix can be reproduced with appropriate substitutions in the results of ref. [38], taking into account the different renormalization conditions on the charged-Higgs mass. We thank Christoph Bobeth for performing this useful comparison.

References

- [1] S. L. Glashow and S. Weinberg, Phys. Rev. D **15** (1977) 1958.
- [2] R. S. Chivukula and H. Georgi, Phys. Lett. B **188** (1987) 99;
G. D'Ambrosio, G. F. Giudice, G. Isidori and A. Strumia, Nucl. Phys. B **645** (2002) 155 [arXiv:hep-ph/0207036].
- [3] A. V. Manohar and M. B. Wise, Phys. Rev. D **74** (2006) 035009 [arXiv:hep-ph/0606172].
- [4] F. J. Botella, G. C. Branco and M. N. Rebelo, arXiv:0911.1753 [hep-ph].
- [5] A. Pich and P. Tuzon, Phys. Rev. D **80** (2009) 091702 [arXiv:0908.1554 [hep-ph]].
- [6] G. C. Branco, W. Grimus and L. Lavoura, Phys. Lett. B **380** (1996) 119 [arXiv:hep-ph/9601383].
- [7] M. I. Gresham and M. B. Wise, Phys. Rev. D **76** (2007) 075003 [arXiv:0706.0909 [hep-ph]].
- [8] A. Idilbi, C. Kim and T. Mehen, Phys. Rev. D **79** (2009) 114016 [arXiv:0903.3668 [hep-ph]].
- [9] R. Bonciani, G. Degrossi and A. Vicini, JHEP **0711** (2007) 095 [arXiv:0709.4227 [hep-ph]].
- [10] C. P. Burgess, M. Trott and S. Zuberi, JHEP **0909** (2009) 082 [arXiv:0907.2696 [hep-ph]].
- [11] H. Flacher, M. Goebel, J. Haller, A. Hocker, K. Moenig and J. Stelzer, Eur. Phys. J. C **60** (2009) 543 [arXiv:0811.0009 [hep-ph]].
- [12] O. Deschamps, S. Descotes-Genon, S. Monteil, V. Niess, S. T'Jampens and V. Tisserand, arXiv:0907.5135 [hep-ph].
- [13] M. Bona *et al.* [UTfit Collaboration], Phys. Lett. B **687** (2010) 61 [arXiv:0908.3470 [hep-ph]].
- [14] J. P. Idarraga, R. Martinez, J. A. Rodriguez and N. Poveda, Braz. J. Phys. **38** (2008) 531.
- [15] J. L. Diaz-Cruz, J. Hernandez-Sanchez, S. Moretti, R. Noriega-Papaqui and A. Rosado, Phys. Rev. D **79** (2009) 095025 [arXiv:0902.4490 [hep-ph]].
- [16] F. Mahmoudi and O. Stal, Phys. Rev. D **81** (2010) 035016 [arXiv:0907.1791 [hep-ph]].
- [17] A. Denner, R. J. Guth, W. Hollik and J. H. Kuhn, Z. Phys. C **51** (1991) 695.
- [18] K. G. Chetyrkin, M. Misiak and M. Munz, Phys. Lett. B **400** (1997) 206 [Erratum-ibid. B **425** (1998) 414] [arXiv:hep-ph/9612313], and references therein.
- [19] M. Ciuchini, G. Degrossi, P. Gambino and G. F. Giudice, Nucl. Phys. B **527** (1998) 21 [hep-ph/9710335].
- [20] P. Ciafaloni, A. Romanino and A. Strumia, Nucl. Phys. B **524** (1998) 361 [arXiv:hep-ph/9710312].

- [21] F. Borzumati and C. Greub, Phys. Rev. D **58** (1998) 074004 [arXiv:hep-ph/9802391], Phys. Rev. D **59** (1999) 057501. [arXiv:hep-ph/9809438].
- [22] H. E. Haber and H. E. Logan, Phys. Rev. D **62** (2000) 015011 [arXiv:hep-ph/9909335].
- [23] S. Davidson and H. E. Haber, Phys. Rev. D **72** (2005) 035004 [Erratum-ibid. D **72** (2005) 099902] [arXiv:hep-ph/0504050].
- [24] T. P. Cheng and M. Sher, Phys. Rev. D **35** (1987) 3484.
- [25] K. G. Chetyrkin, J. H. Kuhn and A. Kwiatkowski, arXiv:hep-ph/9503396.
- [26] J. Fleischer, O. V. Tarasov, F. Jegerlehner and P. Raczka, Phys. Lett. B **293** (1992) 437; G. Buchalla and A. J. Buras, Nucl. Phys. B **398** (1993) 285; G. Degrassi, Nucl. Phys. B **407** (1993) 271 [arXiv:hep-ph/9302288].
- [27] R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci and A. Vicere, Phys. Lett. B **288**, 95 (1992) [Erratum-ibid. B **312**, 511 (1993)] [arXiv:hep-ph/9205238].
- [28] [Tevatron Electroweak Working Group and CDF Collaboration and D0 Collab], arXiv:0903.2503 [hep-ex].
- [29] [ALEPH Collaboration and DELPHI Collaboration and L3 Collaboration and], Phys. Rept. **427** (2006) 257 [arXiv:hep-ex/0509008].
- [30] G. Degrassi, P. Gambino and P. Slavich, Comput. Phys. Commun. **179** (2008) 759 [arXiv:0712.3265 [hep-ph]].
- [31] G. Degrassi, P. Gambino and P. Slavich, Phys. Lett. B **635** (2006) 335 [arXiv:hep-ph/0601135].
- [32] P. Gambino and M. Misiak, Nucl. Phys. B **611** (2001) 338 [arXiv:hep-ph/0104034].
- [33] M. Misiak *et al.*, Phys. Rev. Lett. **98** (2007) 022002 [arXiv:hep-ph/0609232].
- [34] P. Gambino and P. Giordano, Phys. Lett. B **669** (2008) 69 [arXiv:0805.0271 [hep-ph]].
- [35] E. Barberio *et al.* [Heavy Flavor Averaging Group], arXiv:0808.1297 [hep-ex].
- [36] P. Gambino, U. Haisch and M. Misiak, Phys. Rev. Lett. **94** (2005) 061803 [arXiv:hep-ph/0410155].
- [37] A. L. Kagan and M. Neubert, Phys. Lett. B **539** (2002) 227 [arXiv:hep-ph/0110078].
- [38] C. Bobeth, M. Misiak and J. Urban, Nucl. Phys. B **567** (2000) 153 [arXiv:hep-ph/9904413].